

Review Article

Non Surgical Management of Temporomandibular Joint Disorders

Shivangi Pandey, Shitij Srivastava, Abhinav Shekhar, Love Kumar Bhatia

Department of Prosthodontics and Crown Bridge, Sardar Patel Post Graduate Institute of Dental & Medical Sciences, Lucknow (Uttar Pradesh)

ABSTRACT:

Temporomandibular Joint Disorders (TMDs) are multifactorial conditions affecting the temporomandibular joint (TMJ), often presenting as pain, joint sounds, limited jaw movement, and associated myofascial symptoms. This article explores the pathophysiology, diagnosis briefly, and non-surgical therapeutic modalities in detail employed in the management of TMD. Emphasis is placed on conservative and minimally invasive interventions that aim to alleviate pain, restore function, and prevent progression.

Anatomically, the TMJ's unique structure allows both rotational and translational movements, making it vulnerable to mechanical, inflammatory, and neuromuscular disruptions. TMDs may result from trauma, arthritis, bruxism, or psychological stress. Diagnosis involves clinical evaluation, imaging, and classification based on joint or muscular origin.

Pharmacological management includes NSAIDs, antidepressants, and anticonvulsants to modulate inflammation and pain pathways. Non-invasive modalities such as physical therapy, low-level laser therapy (LLLT), ultrasound, and electrical stimulation are utilized to improve joint function and reduce muscular tension. Occlusal splints offer stabilization and occlusal correction in cases of bruxism. Minimally invasive techniques like botulinum toxin (BTX) and hyaluronic acid (HA) injections have demonstrated efficacy in reducing hyperactivity of masticatory muscles and improving joint lubrication, respectively. Arthrocentesis serves as a therapeutic joint lavage to reduce intra-articular inflammation and adhesions.

Overall, non-surgical management strategies provide significant symptomatic relief in most TMD cases, with individualized treatment plans optimizing patient outcomes. The shift from surgical to biopsychosocial and multidisciplinary approaches marks a paradigm evolution in TMD management, highlighting the importance of integrated, evidence-based care.

KEY WORDS: Temporomandibular Joint, regenerative medicine, intra articular injection, pharmacotherapy, hyaluronic acid

Address for correspondence : Dr Shivangi Pandey, Post Graduate Student, Department of Prosthodontics and Crown Bridge, Sardar Patel Post Graduate Institute of Dental & Medical Sciences, Lucknow-226029 (Uttar Pradesh))- India. E-mail: shivangipandey23feb@gmail.com **Submitted:** 23.12.2025, **Accepted:** 07.06.2026, **Published:** 24.06.2026

INTRODUCTION:

The temporomandibular joint (TMJ) connects the mandible to the temporal bone, facilitating essential functions including speech, mastication, and yawning. The TMJ is unique in its capacity for both rotational and translational movements, enabled by an articular

disc that separates the joint compartments. Supporting structures, including muscles, ligaments, and tendons, ensure joint stability and controlled movement. The complex nature and frequent use of the TMJ predispose it to temporomandibular disorders (TMDs), which can manifest as pain, stiffness, joint sounds, and limited jaw movement. TMDs are multifactorial, with common

Access this article online

Quick Response Code:



Website:

www.pjsr.org

DOI:

<https://doi.org/10.5281/zenodo.20841150>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial ShareALike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: editor.pjsr@peoplesuniversity.edu.in

How to cite this article: Pandey S, Srivastava S, Shekhar A, Bhatia LK. Non Surgical Management of Temporomandibular Joint Disorders. PJSR. 2026;19(1):62e.

etiologies including trauma, arthritis, bruxism, and stress. Management of TMDs aims to alleviate pain, reduce joint sounds, and restore normal function. Optimal TMJ health is crucial for overall comfort and proper jaw function.^[1]

HISTORY:

Evolution of TMD Understanding and Treatment:

Historically, malocclusion was considered a primary cause of TMDs. However, research in the 1990s indicated a minimal association, leading to the understanding that orthodontic treatment is not an effective intervention for TMDs. The early 2000s saw the introduction of invasive and surgical treatments, but the treatment paradigm has since shifted towards a biopsychosocial model. Current common treatments include nonsteroidal anti-inflammatory drugs (NSAIDs), muscle relaxants, occlusal splints for bruxism, and physical therapy. Psychological interventions such as relaxation techniques and biofeedback are used to address contributing factors like anxiety. In select cases, corticosteroid or botulinum toxin injections may be indicated, with surgery reserved for severe joint damage.^[2]

TEMPOROMANDIBULAR JOINT DISORDERS:

CLASSIFICATION: The American Academy of Orofacial Pain (AAOP) classification system, integrated with the International Headache Society's^[3] diagnostic criteria, categorizes TMDs into:

1. Temporomandibular joint disorders, including:

- o Deviation in form (articular surface defects, disc thinning)
- o Disc displacement (with or without reduction)
- o Displacement of disc-condyle complex (hypermobility, dislocation)
- o Inflammatory conditions (capsulitis, synovitis, retrodiscitis)
- o Degenerative diseases (osteoarthrosis, polyarthritis, osteoarthritis)
- o Ankylosis (fibrous, bony)

2. Masticatory muscle disorders:

- o Acute (myositis, reflex muscle splinting, muscle spasm)
- o Chronic (myofascial pain, muscle contracture, hypertrophy, myalgia)

3. Congenital and developmental disorders (condylar hyperplasia, hypoplasia, aplasia, neoplasms, fractures)

Okeson's classification further categorizes TMDs into derangements of the condyle-disc complex, structural incompatibilities of the articular surfaces, and inflammatory joint disorders.

DISORDERS: Masticatory muscle disorders are characterized by pain during function and tenderness upon palpation, often with restricted jaw movement.

Specific TMJ disorders include:

- Derangement of the Condyle-Disc Complex: Abnormal condyle-disc rotation due to ligament issues or disc thinning.
- Disc Displacement:
 - o With Reduction: Disc displacement during mouth closure, with repositioning upon opening, often causing joint sounds.
 - o Without Reduction: Disc fails to return to its normal position, causing limited opening, pain, and locking.
- Structural Incompatibility: Alterations in articular surfaces that impede joint function.
 - o Adhesions: Lack of lubrication between disc and condyle or fossa, causing brief restrictions.
 - o Subluxation: Condyle shifts out of position during mouth opening.
 - o Luxation (Dislocation): Condyle dislocates anterior to the articular eminence, preventing closure.
- Inflammatory Disorders: Joint inflammation, including synovitis, capsulitis, retrodiscitis, and arthritis.
 - o Arthritis: Joint inflammation, often with changes in occlusion.
 - o Osteoarthritis: Degenerative condition with pain, disc dislocation, and bone erosion.
- Chronic Mandibular Hypomobility: Long-term limited jaw movement, often painless, due to ankylosis, muscle contracture, or obstruction.
- Growth Disorders: Anomalies in TMJ bone or muscle growth.^[4]

SIGNS AND SYMPTOMS OF TMD: Common signs and symptoms of TMD include jaw pain, joint sounds, limited jaw movement, headaches, facial muscle tenderness, ear symptoms (tinnitus, fullness), painful chewing, neck and shoulder pain.^[4]

PHARMACOTHERAPY:

NSAIDs are commonly used medications for pain and inflammation management. They work by inhibiting cyclooxygenase (COX) enzymes, which are essential in prostaglandin synthesis. Prostaglandins mediate pain and inflammation, and by reducing their production, NSAIDs help alleviate symptoms. Pharmacological approaches for TMD primarily target pain and inflammation.

Pharmacological approaches for TMD primarily target pain and inflammation.

- **Non-Steroidal Anti-Inflammatory Drugs (NSAIDs):** Commonly employed for pain and inflammation, NSAIDs inhibit COX enzymes, thereby reducing prostaglandin synthesis, key mediators of these processes. They inhibit both COX-1, involved in physiological functions, and COX-2, induced during inflammation. Inhibition of COX-2 is the primary mechanism for their anti-inflammatory and analgesic effects in TMDs, alleviating acute and chronic symptoms. Commonly used NSAIDs include ibuprofen, naproxen, diclofenac, and the selective COX-2 inhibitor celecoxib, each with varying COX selectivity and safety profiles. While effective for pain relief in TMD, potential gastrointestinal, cardiovascular, and renal side effects, particularly with prolonged use, necessitate careful consideration.^[5]
- **Antidepressants:** Primarily indicated for mood disorders, antidepressants also demonstrate efficacy in managing chronic pain, including TMD. They modulate pain pathways and central sensitization. Selective serotonin reuptake inhibitors (SSRIs) enhance serotonin availability. Serotonin-norepinephrine reuptake inhibitors (SNRIs) target both neurotransmitters, proving beneficial for neuropathic and musculoskeletal pain. Tricyclic antidepressants (TCAs) block serotonin and norepinephrine reuptake, exhibiting sedative and analgesic properties. Atypical antidepressants possess diverse mechanisms affecting pain perception. In TMD, antidepressants modulate central pain pathways, relieve muscle spasms (TCAs, SNRIs), improve sleep (sedative antidepressants), and address comorbid psychological factors. Clinical evidence supports their role in TMD pain reduction, although side effects require monitoring.^[6]

- **Anticonvulsants:** Primarily used for epilepsy, anticonvulsants are increasingly utilized for chronic pain management, including TMD. They modulate neuronal excitability, proving effective for neuropathic pain and central sensitization. Mechanisms include sodium channel inhibition (e.g., carbamazepine), calcium channel modulation (e.g., gabapentin, pregabalin), enhancement of GABAergic inhibition (e.g., valproate), and glutamate modulation. In TMD, they offer relief from neuropathic pain, reduce central sensitization, and can be adjunctive in refractory cases. Sedative effects may also improve sleep and anxiety. While studies show pain reduction, potential side effects necessitate careful consideration.^[7]

PHYSICAL THERAPY:-

LASER THERAPY: Utilizing red or infrared electromagnetic radiation, LLLT is employed for localized musculoskeletal pain, including TMD. Its bio-stimulatory, regenerative, analgesic, and anti-inflammatory effects may involve endogenous opioid release, enhanced glucocorticoid excretion, improved microcirculation, reduced nerve permeability, and increased ATP production.^[8] In TMD, LLLT offers a non-invasive adjunct to other therapies, potentially improving pain tolerance, blood flow, cellular metabolism, and mouth opening.^[9,10,11]

SPLINT THERAPY: A widely used non-invasive treatment for TMD, splints stabilize the TMJ, reduce strain, improve jaw alignment, and prevent joint damage, particularly in cases of bruxism and malocclusion. Stabilization splints maintain a neutral jaw position, while repositioning splints aim to realign the condyle.^[12] Anterior bite plane splints target bruxism. Custom-made splints offer optimal fit and effectiveness. Splint therapy provides symptomatic relief and protects teeth, but may involve initial discomfort.^[13] Table 1 shows different types of splints used in TMJ disorders.

Benefits and Challenges of Splint Therapy: Splint therapy provides non-invasive relief, restores joint function, and protects teeth from bruxism. However, initial discomfort, adaptation periods, and potential complications like increased salivation or difficulty speaking may occur.^[13]

ULTRASOUND AND ELECTRIC STIMULATION:

These non-invasive treatments are used for TMD. Ultrasound utilizes sound waves for thermal and

Table 1: Types of Splints Used in TMJ Disorders:

Stabilization Splints	These splints cover the teeth to hold the jaw in a neutral position. They are made of hard acrylic and help prevent tooth contact during bruxism, reducing muscle fatigue and joint strain.
Repositioning Splints	Used to realign the jaw, these splints are designed to reposition the condyle within the glenoid fossa, addressing issues like malocclusion or disc displacement. They are often worn longer to gradually change occlusion
Anterior Bite Plane Splints	These splints target bruxism and excessive clenching by isolating posterior teeth and preventing contact. They reduce muscle tension and jaw pressure
Soft Splints	Made from flexible material, soft splints offer temporary relief by cushioning the teeth and reducing pressure on the TMJ, but are less effective long-term.
Custom-Made vs. Over-the-Counter Splints	Custom-made splints are tailored to the patient for optimal fit and effectiveness, while over-the-counter splints offer a more general, temporary solution.
Hybrid Splints	These combine features of stabilization and repositioning splints, addressing both misalignment and bruxism

non-thermal effects, improving blood flow and promoting tissue healing. Electrical stimulation aims to alleviate pain and restore function. Both are often used adjunctively.^[14]

PHYSICAL EXERCISE IN TMD MANAGEMENT:

Exercise plays a key role in managing TMD by alleviating pain, improving jaw mobility, and preventing long-term complications. Regular, controlled exercises restore normal jaw movement, reduce muscle tension, and prevent jaw locking. Strengthening the muscles surrounding the TMJ helps reduce stress, inflammation, and discomfort.

Types of Exercises for TMD:

1. Range-of-Motion Exercises:

- *Opening and Closing:* Gently open and close the mouth to improve jaw mobility.
- *Side-to-Side:* Move the jaw laterally to increase lateral mobility.
- *Protrusion and Retrusion:* Move the jaw forward and backward to enhance flexibility.

2. Strengthening Exercises:

- *Jaw Resistance:* Apply resistance to the chin while opening the mouth to strengthen masticatory muscles.

- *Isometric Exercises:* Contract jaw muscles without joint movement to build strength.

3. Relaxation and Stretching Exercises:

- *Progressive Muscle Relaxation:* Relax jaw muscles to reduce tension.
- *Stretching:* Stretch jaw muscles gently to alleviate tightness.
- *Neck and Postural Exercises:* Improve posture and relieve tension in related muscles.^[14]

MINIMALLY INVASIVE TREATMENTS:-

MUSCLE INJECTION WITH BTX: *Botulinum Toxin (BTX) in the Treatment of Temporomandibular Disorders:* Botulinum toxin, a neurotoxin produced by *Clostridium botulinum*, has become a transformative therapeutic agent in medical and dental fields. Initially recognized for its role in cosmetic treatments, BTX is increasingly utilized for managing neuromuscular conditions, including TMDs. By targeting hyperactive muscles and modulating pain pathways, BTX offers a promising treatment modality for TMDs, especially for refractory cases.

Overview of Botulinum Toxin:

BTX inhibits acetylcholine release at the neuromuscular junction, leading to temporary muscle paralysis. BTX-A (e.g., onabotulinumtoxin A) and BTX-B (e.g., rimabotulinumtoxin B) are commonly

used for managing TMDs by reducing muscle hyperactivity and modulating pain.

Temporomandibular Disorders: Pathophysiology and Treatment Challenges:

TMDs include muscle-related pain (myofascial pain), joint-related issues, and mixed disorders. BTX therapy is particularly beneficial for refractory cases, especially when conservative treatments fail^[15]

Mechanism of Action of BTX in TMD **BTX alleviates TMD symptoms through:**

- **Muscle Relaxation:** Reduces muscle contractions in hyperactive muscles.
- **Reduction of Peripheral Pain:** Lowers pain-mediating substances.
- **Central Modulation of Pain:** Reduces hyperalgesia and allodynia.

Improvement of Bruxism: Weakens jaw-closing muscles, reducing clenching and grinding.

Mechanism of Action of BTX in TMD

BTX alleviates symptoms of TMD through multiple mechanisms:

1. **Muscle Relaxation**
 - By inhibiting acetylcholine release, BTX reduces muscle contractions in hyperactive or hypertrophic masticatory muscles, alleviating pain and improving jaw function.
2. **Reduction of Peripheral Pain**
 - BTX decreases the release of pain-mediating substances such as substance P, glutamate, and calcitonin gene-related peptide (CGRP) from peripheral nerve endings, leading to reduced nociception.
3. **Central Modulation of Pain**
 - Evidence suggests that BTX indirectly modulates central pain processing, reducing hyperalgesia and allodynia associated with chronic TMD.
4. **Improvement of Bruxism and Parafunctional Habits**
 - BTX weakens jaw-closing muscles such as the masseter and temporalis, reducing the intensity of clenching and grinding behaviors^[16]

Indications for BTX in TMD:

BTX is indicated for various TMD presentations, particularly in cases where traditional treatments have proven insufficient:

- Myofascial Pain Dysfunction Syndrome

- Severe Bruxism
- Hypertrophic Masticatory Muscles
- Chronic Migraine or Headache Associated with TMD
- Post-Surgical or Refractory TMJ Pain

Clinical Application

- **Masseter:** For bruxism, hypertrophy, and general myofascial pain.
- **Temporalis:** For tension-related headaches and temporal muscle pain.
- **Lateral Pterygoid:** For disc displacement or TMJ-related dysfunction.

Dosing varies depending on muscle size, severity of symptoms, and individual response. BTX effects typically manifest within 3–7 days, peak at 2–4 weeks, and last 3–6 months, necessitating repeat injections for sustained benefit^[17]

Safety and Side Effects:

BTX is generally well-tolerated, but potential side effects must be considered:

1. **Local Effects**
 - Temporary weakness of adjacent muscles, difficulty chewing, or asymmetry in facial expressions.
2. **Systemic Effects**
 - Rare cases of distant toxin spread may result in flu-like symptoms or generalized muscle weakness.
3. **Reversibility**
 - Effects are temporary, requiring periodic re-administration, which may be burdensome for some patients.
4. **Cost**
 - High costs may limit accessibility, particularly for patients requiring repeated treatments.

Future Directions:

Emerging research is exploring novel formulations and delivery methods for BTX to enhance its efficacy and duration of action. Studies are also investigating its combined use with other therapies, such as physical therapy or pharmacological agents, to optimize outcomes. Advances in imaging-guided injection techniques aim to improve precision and minimize side effects.^[18]

Hyaluronic Acid Injections in Temporomandibular Disorders:

Hyaluronic acid (HA) is a naturally occurring glycosaminoglycan found in the extracellular matrix

(ECM) and synovial fluid of diarthrodial joints, including the TMJ. It plays a crucial role in maintaining the structural integrity and viscosity of synovial fluid, contributing to its lubricating properties. Synthesized by synoviocytes, HA is essential for joint homeostasis. Its molecular structure consists of repeating disaccharide units of D-glucuronic acid and N-acetyl-D-glucosamine linked by β -1,4 and β -1,3 glycosidic bonds, resulting in a high-molecular-weight polymer (approximately 6000-7000 kDa).^[19]

Mechanism of Action in TMDs: When injected into the TMJ, HA exerts several therapeutic effects:

- **Restoration of Joint Lubrication:** HA replenishes the viscoelastic properties of synovial fluid, reducing friction between articular surfaces and improving joint movement.
- **Reduction of Inflammation:** By decreasing the production of inflammatory mediators and matrix metalloproteinases, HA helps alleviate synovitis and joint pain.
- **Cartilage Protection and Repair:** HA can stimulate chondrocytes and inhibit cartilage degradation, potentially slowing the progression of degenerative changes within the TMJ.
- **Flushing of Debris:** When combined with arthrocentesis, HA injections can aid in the removal of inflammatory debris and degraded cartilage particles from the joint space.
- **Improved Tissue Healing:** HA can enhance the proliferation and migration of fibroblasts and synoviocytes, promoting tissue repair within the joint capsule.

Clinical Indications for HA in TMD: HA injections are particularly beneficial for:

- **Internal Derangements:** Disorders involving disc displacement without reduction, leading to pain and limited mandibular movement.
- **TMJ Osteoarthritis:** Degenerative joint disease characterized by cartilage breakdown, subchondral bone changes, and joint space narrowing.
- **Synovitis or Capsulitis:** Inflammatory conditions causing pain, swelling, and stiffness in the TMJ.
- **Post-Surgical Rehabilitation:** Following TMJ surgeries, HA injections can aid in restoring joint function and reducing inflammation.
- **Refractory TMD Symptoms:** Patients

unresponsive to conservative therapies may benefit from HA as an adjunctive treatment.^[20]

Administration of HA in TMD: HA injections are performed under aseptic conditions, often guided by ultrasound or fluoroscopy for precise intra-articular delivery. Key steps include:

- **Pre-Injection Assessment:** A comprehensive evaluation of TMJ function, imaging studies, and patient-specific factors is essential.
- **Injection Protocol:** HA is injected directly into the joint space, sometimes following arthrocentesis to flush the joint. Dosages and formulations vary based on the severity of the disorder.
- **Post-Injection Care:** Patients are typically advised to avoid excessive jaw movements and may benefit from follow-up physical therapy or other supportive treatments.^[20]

ARTHROCENTESIS: Arthrocentesis is a minimally invasive procedure involving the lavage or flushing of the TMJ with sterile fluids to reduce inflammation and improve joint mobility.^[21]

Advantages of Arthrocentesis:

- **Minimally Invasive:** Involves no surgical incisions, resulting in a shorter recovery period.
- **Immediate Relief:** Can provide relatively rapid reduction in pain and improvement in joint mobility.
- **Reduction of Inflammation:** The flushing action helps to wash out inflammatory mediators from the joint space.

Disadvantages and Limitations:

- **Limited Indication for Advanced Pathology:** May not be suitable for severe osteoarthritis or intra-articular tumors.
- **Temporary Relief:** The effects of arthrocentesis may not be long-lasting for all patients.
- **Technical Challenges:** Requires a skilled practitioner for proper cannula placement and joint lavage.
- **Lack of Structural Repair:** Does not address underlying structural damage within the joint.
- **Variability in Outcomes:** The success of arthrocentesis can vary depending on individual patient factors and the specific TMD pathology.^[22]

CONCLUSION:

Management of TMD necessitates a comprehensive and individualized approach, encompassing both conservative and minimally invasive treatments tailored to the specific needs of each patient. The primary objectives are to alleviate pain, restore optimal joint function, and improve the patient's overall quality of life. Initial management often involves conservative strategies such as NSAIDs, muscle relaxants, and physical therapy. These therapies may include targeted jaw exercises, postural correction techniques, and the use of occlusal appliances (night guards) to manage bruxism. When conservative measures prove insufficient in providing adequate relief, minimally invasive advanced treatments like BTX or HA injections may be considered to reduce muscle hyperactivity and inflammation within the joint, respectively. Arthrocentesis, a joint lavage procedure, can also be employed to reduce inflammation and improve joint mobility. Surgical options are typically reserved for severe cases involving significant structural damage to the TMJ. Dietary modifications, such as avoiding hard or chewy foods to minimize strain on the jaw, alongside improving sleep hygiene and addressing nocturnal bruxism, are crucial components of symptom management. Effective TMD treatment is inherently individualized and often requires ongoing collaboration among dentists, physical therapists, and other healthcare professionals to achieve the best possible outcomes. With appropriate and timely management, the majority of patients with TMD can experience significant relief from their symptoms and a substantial improvement in their quality of life.

Financial Support and Sponsorship

Nil.

Conflicts of Interest

There are no conflicts of interest.

REFERENCES:

1. Dolwick MF. The role of temporomandibular joint surgery in the treatment of patients with internal derangement. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1997;83:150–5.
2. McCarty WL, Farrar WB. Surgery for internal derangements of the temporomandibular joint. *J Prosthet Dent.* 1979;42:191–6.
3. Yakkaphan P, Smith JG, Chana P, Tan HL, Ravindranath PT, Lambru G, Renton T. Temporomandibular Disorders and Fibromyalgia Prevalence: A Systematic Review and Meta-Analysis. *J Oral Facial Pain Headache.* 2023;17;37(3):177–193. Doi: 10.11607/ofph.3260. PMID: 37975782; PMCID:PMC10664703.
4. Sipahi A, Satilmis T, Basa S. Comparative study in patients with symptomatic internal derangements of the temporomandibular joint: analgesic outcomes of arthrocentesis with or without intraarticular morphine and tramadol. *Br J Oral Maxillofac Surg.* 2015;53:316–320.
5. Herman CR, Schiffman EL, Look JO, Rindal DB. The effectiveness of adding pharmacologic treatment with clonazepam or cyclobenzaprine to patient education and self-care for the treatment of jaw pain upon awakening: a randomized clinical trial. *J Orofac Pain.* 2002;16:64–70.
6. Rizzatti-Barbosa CM, Nogueira MT, de Andrade ED, Ambrosano GM, de Barbosa JR. Clinical evaluation of amitriptyline for the control of chronic pain caused by temporomandibular joint disorders. *Cranio.* 2003;21:221–225.
7. Hersh EV, Balasubramaniam R, Pinto A. Pharmacologic management of temporomandibular disorders. *Oral Maxillofac Surg Clin North Am.* 2008;20:197–210.
8. Ownby SL, Fortuno LV, Au AY, Grzanna MW, Rashmir-Raven AM, Frondoza CG. Expression of pro-inflammatory mediators is inhibited by an avocado/soybean unsaponifiables and epigallocatechin gallate combination. *J Inflamm (Lond)* 2014;11:8.
9. Ahrari F, Madani AS, Ghafouri ZS, Tunér J. The efficacy of low level laser therapy for the treatment of myogenous temporomandibular joint disorder. *Lasers Med Sci.* 2014;29:551–557.
10. Coombe AR, Ho CT, Darendeliler MA, Hunter N, Philips JR, Chapple CC, et al. The effects of low level laser irradiation on osteoblastic cells. *Clin Orthod Res.* 2001;4:3–14.
11. Pinheiro AL, Cavalcanti ET, Pinheiro TI, Alves MJ, Miranda ER, De Quevedo AS, et al. Low-level laser therapy is an important tool to treat disorders of the maxillofacial region. *J Clin Laser Med Surg.* 1998;16:223–226.
12. Esposito CJ, Veal SJ, Farman AG. Alleviation of myofascial pain with ultrasonic therapy. *J Prosthet Dent.* 1984;51:106–108.
13. Gray RJ, Davies SJ. Occlusal splints and temporomandibular disorders: why, when, how? *Dent Update.* 2001;28:194–199.
14. Sato S, Nasu F, Motegi K. Analysis of post-treatment electromyographs in patients with non-reducing disc displacement of the temporomandibular joint. *J Oral Rehabil.* 2002;29:1126–1130.
15. Arinci A, Güven E, Yazar M, Başaran K, Keklik B. Effect of injection of botulinum toxin on lateral pterygoid muscle used together with the arthroscopy in patients with anterior disk displacement of the temporomandibular joint. *Kulak Burun Bogaz Ihtis Derg.* 2009;19:122–129.
16. Sidebottom AJ, Patel AA, Amin J. Botulinum injection for the management of myofascial pain in the

- masticatory muscles. A prospective outcome study. *Br J Oral Maxillofac Surg*. 2013;51:199–205.
17. Park SY, Park YW, Ji YJ, Park SW, Kim SG. Effects of a botulinum toxin type A injection on the masseter muscle: an animal model study. *MaxillofacPlastReconstr Surg*. 2015;37:10.
 18. Song PC, Schwartz J, Blitzler A. The emerging role of botulinum toxin in the treatment of temporomandibular disorders. *Oral Dis*. 2007;13:253–260.
 19. Monje-Gil F, Nitzan D, González-García R. Temporomandibular joint arthrocentesis. Review of the literature. *Med Oral Patol Oral Cir Bucal*. 2012;17:e575–e581.
 20. Özgür Yİs M, Erbaş D. Effect of arthrocentesis and sodium hyaluronate injection on nitrite, nitrate, and thiobarbituric acid-reactive substance levels in the synovial fluid. *Oral Surg Oral Med Oral Pathol Oral RadiolEndod*. 2000;89:686–690.
 21. Basterzi Y, Sari A, Demirkan F, Unal S, Arslan E. Intraarticular hyaluronic acid injection for the treatment of reducing and nonreducing disc displacement of the temporomandibular joint. *Ann Plast Surg*. 2009;62:265–267.]
 22. Yeung RWK, Chow RLK, Samman N, Chiu K. Short-term therapeutic outcome of intra-articular high molecular weight hyaluronic acid injection for nonreducing disc displacement of the temporomandibular joint. *Oral Surg Oral Med Oral Pathol Oral RadiolEndod*. 2006;102:453–461.